



**Stantec**

**Stantec Consulting Ltd.**  
40 Highfield Park Drive, Suite 102  
Dartmouth NS B3A 0A3  
Tel: (902) 468-7777  
Fax: (902) 468-9009

December 10, 2012  
File: 121510982

Clayton Developments Limited  
255 Lacewood Drive, Suite 100C  
Halifax, NS B3M 4G2

**Attention:** **Mike Hanusiak, Sr. Vice President, General Manager**

Dear Mr. Hanusiak:

**Reference:** **Water Quality Monitoring Results for Russell Lake – November 2012 Sampling Event**

To monitor the effects of development, water quality monitoring has been undertaken on a seasonal basis since April of 2005 at four stations in Russell Lake (Figure 1 in Attachment). Monitoring events include one spring, two summer and one fall sampling event each year. Water samples are not collected in winter. Sampling results have been provided in previous letter reports to you, with a summary of the sampling up to June 27, 2012 presented in the report dated August 12, 2012. The present report builds on the results to date while focusing on the most current fall sampling event on November 7, 2012, with special attention provided to previous fall sampling events.

The weather during the sampling period on November 7, 2012 was mainly clear with an air temperature of -4 °C to 2 °C. A trace amount of rain occurred within 48 hours prior to the sampling event, with 0.5 mm of rain measured at the Halifax International Airport on November 5. Hourly mean wind speeds were up to 29 km/h and blew primarily from the north for 48 hours leading up to the date of sample collection. The mean hourly wind speeds were between 13 and 29 km/h on the sampling date, higher than the previous 48 hours.

Sampling was conducted between 9:30 and 11:40 on November 7, 2012. The following conditions were observed at the four Russell Lake monitoring stations (i.e., In-Lake, Outlet, South Inlet and North Inlet sampling stations):

- In Russell Lake, the water appeared clear to slightly tea-stained. Three ducks were observed in the vicinity of the boat slipway and two ducks observed near the outlet.
- At the Outlet station of Russell Lake, the water appeared clear. The water level at the Outlet station appeared similar to other November events. In-stream vegetation was present, though beginning to decay along the banks. Two ducks were observed approximately 4 m upstream from the sampling location.
- Water flowing in the South Inlet station appeared slightly tea-stained. The South Inlet station contained a water level similar to previous November sampling events.
- The water appeared clear at the time of sample collection from the North Inlet station. The grating within the culvert at this station contained garbage. Algal growth was minimal but present on the substrate within the North Inlet dissipation pool.

Tables 1 to 6 below present statistical summaries of analytical results from April 2005 to November 2012 for key water quality parameters (total suspended solids (TSS), turbidity, pH, conductivity, sodium and chloride, total phosphorus (TP), Chlorophyll a (Chl a), and faecal coliforms) at each of the monitoring stations. The seasonal trends for some of the key water quality parameters are presented graphically in Figures 2 to 8 below. Where applicable, the water quality data were compared to the Canadian Council for the Ministers of the Environment Guidelines for the Protection of Freshwater Aquatic Life (CCME FAL).

December 10, 2012

Mike Hanusiak, Sr. Vice President, General Manager

Page 2 of 6

**Reference: Water Quality Monitoring Results for Russell Lake – November 2012 Sampling Event****Water Clarity**

TSS concentrations (Figure 2 in Attachment and Table 1 below) were not detected and below the reportable detection limit of 1 mg/L at all stations during the fall monitoring event. These results are consistent with low TSS concentrations for previous November sampling events.

**Table 1 Russell Lake - TSS Statistics**

Station	November 7, 2012 Results (mg/L)	Fall Season Min (mg/L)	Fall Season Max (mg/L)	Fall Season Median (mg/L)	Fall Season Mean (mg/L)
In-Lake	ND	ND	4	ND	2
Outlet	ND	ND	3	2	2
South Inlet	ND	ND	5	2	2
North Inlet	ND	ND	3	2	2

ND = &lt;1 mg/L TSS

Turbidity levels at the In-Lake, Outlet and North Inlet stations for November 7, 2012 were generally consistent with historical fall median values as presented below in Table 2. Turbidity at the South Inlet was the highest of all fall samples at that station to date and also the highest turbidity value inclusive of all the stations sampled in November 2012.

**Table 2 Russell Lake - Turbidity Statistics**

Station	November 7, 2012 Results (NTU)	Fall Season Min (NTU)	Fall Season Max (NTU)	Fall Season Median (NTU)	Fall Season Mean (NTU)
In-Lake	1.3	0.7	3.6	1.2	1.8
Outlet	1.3	0.3	2.8	1.1	1.4
South Inlet	5.5	0.7	4.5	1.5	1.9
North Inlet	1.3	0.6	10.0	1.4	2.8

**pH**

Measurements of pH collected since April 2005 (Figure 3 in Attachment and Table 3 below) indicate that the water of Russell Lake and its tributaries are historically slightly basic with a median value of 7.5. The pH readings in November 2012 were neutral to slightly basic and ranged from 7.0 at the North Inlet station to 8.2 at the Outlet station. The pH results for the In-Lake and Outlet stations are within the CCME FAL guideline for pH of 6.5 to 9.0 though represent the highest values measured at those stations during the fall season since monitoring began in 2005.

**Table 3 Russell Lake - pH Statistics**

Station	November 7, 2012 Results (Units)	Fall Season Min (Units)	Fall Season Max (Units)	Fall Season Median (Units)	Fall Season Mean (Units)
In-Lake	8.0	5.7	7.8	7.2	7.0
Outlet	8.2	6.8	7.8	7.4	7.4
South Inlet	7.3	5.7	7.6	7.3	7.0
North Inlet	7.0	7.0	8.1	7.5	7.5

December 10, 2012

Mike Hanusiak, Sr. Vice President, General Manager

Page 3 of 6

**Reference: Water Quality Monitoring Results for Russell Lake – November 2012 Sampling Event****Salt Concentrations**

As is typical of urban lakes in watersheds subject to development, Russell Lake has relatively higher salt concentrations than would be expected for a similar lake in a less developed watershed. Conductivity increases with salt concentrations and elevated levels are typically exhibited during spring melt conditions that generally decrease to lower concentrations during the summer monitoring periods. Concentrations tend to be higher at the North Inlet station, where the road network and associated drainage is concentrated and lower at the South Inlet station, which is less developed.

As presented in Table 4 (below) and on Figure 4 (Attachment), the November 2012 conductivity measurements at the In-Lake, Outlet and North Inlet stations were within the range observed since 2005. The conductivity of the South Inlet waters was equal to the previously recorded fall season maximum that occurred in November 2010.

**Table 4 Russell Lake - Conductivity Statistics**

Station	November 7, 2012 Results ( $\mu\text{S}/\text{cm}$ )	Fall Season Min ( $\mu\text{S}/\text{cm}$ )	Fall Season Max ( $\mu\text{S}/\text{cm}$ )	Fall Season Median ( $\mu\text{S}/\text{cm}$ )	Fall Season Mean ( $\mu\text{S}/\text{cm}$ )
In-Lake	480	310	700	480	498
Outlet	480	300	710	480	496
South Inlet	210	120	210	145	158
North Inlet	890	430	1000	770	738

Sodium and chloride are indicators of road salt concentrations in urban lakes and are discussed in that context below. Given that CCME FAL recommends a chloride concentration of less than 120 mg/L for the long-term health of aquatic organisms (guideline of 640 mg/L for short-term exposure), the discussion of salt concentrations below will revolve predominantly around chloride with a brief description of sodium concentrations within Russell Lake.

Sodium (Na) concentrations in November 2012 at the In-Lake, Outlet, South Inlet and North Inlet sampling stations (74 mg/L, 77 mg/L, 23 mg/L and 116 mg/L, respectively) were within their respective ranges observed over the duration of monitoring during previous fall events.

Chloride ions at the In-Lake, Outlet and North Inlet stations (120 mg/L, 120 mg/L and 210 mg/L, respectively) were similarly within the range observed over the duration of monitoring (refer to Table 5 below and Figure 5 in Attachment). Though the concentrations of chloride at these stations were within historic ranges they met or exceeded the CCME FAL long-term guideline value of 120 mg/L. The In-Lake, Outlet and North Inlet stations on the long term have generally exceeded the CCME long-term guideline concentration for chloride, with the North Inlet having the highest median fall chloride concentration of 170 mg/L (see Table 5). The November 2012 South Inlet sample concentration (38 mg/L) was below the CCME long-term guideline, though it did exceed the maximum fall concentration previously observed in 2008 and 2010 (37 mg/L).

December 10, 2012

Mike Hanusiak, Sr. Vice President, General Manager

Page 4 of 6

**Reference: Water Quality Monitoring Results for Russell Lake – November 2012 Sampling Event****Table 5 Russell Lake - Chloride Statistics**

Station	November 7, 2012 Results (mg/L)	Fall Season Min (mg/L)	Fall Season Max (mg/L)	Fall Season Median (mg/L)	Fall Season Mean (mg/L)
In-Lake	120	72	190	94	118
Outlet	120	72	190	93	118
South Inlet	38	20	37	27	28
North Inlet	210	87	250	170	160

**Nutrient Enrichment**

The Chlorophyll a (Chl a) concentration of 2.87 µg/L (based on the acidification technique) at the In-Lake site for November 2012 was within the range of previously reported fall samples (Table 6 below and Figure 6 in Attachment). The fall season results for the In-Lake station from previous sampling years ranged from 1.48 µg/L in 2008 to 18.67 µg/L in 2011. The Outlet sample (2.30 µg/L) also was within the range of previous fall samples which included a low of 1.04 µg/L in 2008 and a high of 17.70 µg/L in 2005.

Total phosphorus (TP) concentrations at the In-Lake station have fluctuated over the fall sampling events, ranging from a low of 2 µg/L in 2010 to a high of 25 µg/L in 2005 (Figure 7 in Attachment and Table 7). The total phosphorus concentration in November 2012 was 4 µg/L (Table 7), which is below the HRM threshold value of 15 µg/L for Russell Lake. It should be noted that the HRM threshold value is intended for comparison to sample results at the In-Lake station only. The Outlet station TP concentration fluctuated similarly to the In-Lake station with a range from 6 µg/L in 2009 to 27 µg/L in 2005. The November 2012 TP concentration of 8 µg/L was slightly higher than the In-Lake station though within the range previously observed (Figure 7) and below the fall median concentration (Table 7).

**Table 6 Russell Lake - Chl a Statistics**

Station	November 7, 2012 Results (µg/L)	Fall Season Min (µg/L)	Fall Season Max (µg/L)	Fall Season Median (µg/L)	Fall Season Mean (µg/L)
In-Lake	2.87	1.48	18.67	6.58	8.40
Outlet	2.30	1.04	17.70	5.16	7.70
South Inlet	0.24	0.13	1.08	0.29	0.38
North Inlet	0.53	0.170	6.60	1.50	2.10

The Chl a concentration for the November 2012 sampling event at the South Inlet station was 0.24 µg/L (Table 6). This result is within the range of previous fall season values, which were between 0.13 µg/L (in 2010) and 1.08 µg/L (in 2006). The TP concentration at the South Inlet station in November 2012 measured 49 µg/L (Table 7), which is the third highest concentration recorded since 2005. Prior TP concentrations ranged from 26 µg/L in 2010 to 110 µg/L in 2006.

The Chl a concentration for the November 2012 sampling event at the North Inlet station was 0.53 µg/L (Table 6 above). This result is within the range of all previous fall season values, which were between 0.170 µg/L (in 2008) and 6.60 µg/L (in 2007). As indicated in Table 7 (below), the TP concentration at the North Inlet station in November 2012 measured 9 µg/L and is within the range of previous fall values for TP concentrations (6 µg/L in 2010 to 27 µg/L in 2011).

December 10, 2012

Mike Hanusiak, Sr. Vice President, General Manager

Page 5 of 6

**Reference: Water Quality Monitoring Results for Russell Lake – November 2012 Sampling Event****Table 7 Russell Lake - TP Statistics**

Station	November 7, 2012 Results (µg/L)	Fall Season Min (µg/L)	Fall Season Max (µg/L)	Fall Season Median (µg/L)	Fall Season Mean (µg/L)
In-Lake	4	2	25	13	12
Outlet	8	6	27	11	12
South Inlet	49	26	110	44	54
North Inlet	9	6	27	10	12

**Bacterial Contamination**

Bacterial contamination in Russell Lake is measured by sampling and testing for the presence of faecal coliforms in the water. In addition to faecal coliform analysis conducted by the laboratory, analysis of *Escherichia coli* (*E. coli*) has also been carried out for sampling events beginning in 2010. *E. coli* are a type of faecal coliform that may be considered a more specific indicator of bacteria found in the intestines of warm-blooded animals and humans and are a strong indicator of recent sewage or animal waste contamination, whereas some faecal bacteria may originate from non-faecal sources. *E. coli* counts were similar for both the In-Lake and Outlet Stations with counts of 20 CFU/100 mL for each. The *E. coli* counts at the South and North Inlet Stations were 30 CFU/100ml and 60 CFU/100ml, respectively. The concentration of faecal coliforms measured at the In-Lake station was 10 CFU/100 mL and 30 CFU/100 mL at the North Inlet station. The samples for the Outlet and South Inlet stations were below the reportable detection limit of 10 CFU/100 mL. These bacterial concentrations are low for fall sampling events when compared to station-specific results from past years (refer to Figure 8 in Attachment).

**Conclusion**

The results of the November 2012 sampling event indicate that total phosphorus (TP) levels at the In-Lake station are below the HRM threshold value of 15 µg/L. The fall TP result at the Outlet station was slightly higher than the concentrations at the In-Lake station. The In-Lake and Outlet Chl a concentrations were the second lowest concentration, for their respective stations, observed during the fall season.

The November 2012 conductivity, sodium (Na) and chloride (Cl) concentrations were within the range of results from previous fall sampling events for the In-Lake, Outlet and North Inlet stations, though they were at or slightly above their respective maximum historic fall values for the South Inlet station.

The pH values from November 2012 at the In-Lake and Outlet stations exceeded the historic maximum fall values, though they were within the CCME FAL guideline for pH of 6.5 to 9.0.

The TSS, Chl a, TP and Fecal coliform concentrations were all within the range of results from previous fall sampling events and generally the November 2012 concentrations were below the respective median values.

December 10, 2012

Mike Hanusiak, Sr. Vice President, General Manager

Page 6 of 6

**Reference: Water Quality Monitoring Results for Russell Lake – November 2012 Sampling Event**

Sincerely,

**STANTEC CONSULTING LTD.**

*ORIGINAL SIGNED BY*

Matt Steeves, B.Sc  
Aquatic Scientist  
Tel: (902) 468-7777  
[matt.steeves@stantec.com](mailto:matt.steeves@stantec.com)

*ORIGINAL SIGNED BY*

Amber Fox, MREM  
Project Manager  
Tel: (902) 468-7777  
[amber.fox@stantec.com](mailto:amber.fox@stantec.com)

cc: Cameron Deacoff, Halifax Regional Municipality

Attachment: Figures 1-8 and Tables 8-11




Figure 1

## Russell Lake Sampling Locations

Map Parameters  
 Projection: UTM-NAD83-Z20  
 Scale 1:9,000  
 Date: May 11, 2007  
 Project No.: NSD19184

0 250 500  
 Metres






Figure 2. Total Suspended Solids at four sites in Russell Lake from April 2005 to November 2012




Figure 3. pH at four sites in Russell Lake from April 2005 to November 2012




Figure 4. Conductivity at four sites in Russell Lake from April 2005 to November 2012




Figure 5. Chloride concentrations at four sites in Russell Lake from April 2005 to November 2012




Figure 6. Chlorophyll a concentrations at four sites in Russell Lake from April 2005 to November 2012




Figure 7. Total Phosphorous concentrations at four sites in Russell Lake from April 2005 to November 2012




Figure 8. Faecal Coliform at four sites in Russell Lake from April 2005 to November 2012

TABLE 8 Surface Water Quality Data for Russell Lake, In-Lake (2005-2012)

Analyte	Units	CCME FWAL	In-Lake																																		
			2005				2006				2007				2008				2009				2010				2011										
			Apr	Jun	Aug	Nov	Mar	Jun	Aug	Oct	Apr	Jun	Aug	Oct	Mar	Jun	Aug	Oct	Apr	Jun	Aug	Oct	Mar	Jun	Aug	Nov	Apr	Jun	Aug	Nov							
<b>General Chemistry</b>																																					
Total Alkalinity (as CaCO <sub>3</sub> )	mg/L		20	17	21	17	15	20	25	26	22	29	27	30	19	22	28	34	27	30	36	35	30	35	36	36	32	31	33	35	27	33	35	36			
Chloride	mg/L		150	87	99	72	91	100	77	79	170	140	120	89	180	250	220	190	270	250	220	150	170	180	160	150	240	170	130	94	170	190	190	120			
Colour	TCU		12	12	8	18	12	14	13	11	9	9	8	12	10	11	15	9	16	9	14	20	14	8	10	15	14	19	22	21	10	7.9	6.2	15			
Hardness (as CaCO <sub>3</sub> )	mg/L		48	34	40	37		43	39	42	59	53	52	46	52	62	64	56	70	65	66	52	54	69	65	58	69	60	53	47	56	63	62	52			
Nitrate + Nitrite (as N)	mg/L	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	0.15	<0.05	<0.05	<0.05	0.2	<0.05	<0.05	<0.05	0.11	<0.05	<0.05	<0.05	0.08	<0.05	<0.05	<0.05	0.095	<0.05	<0.05	<0.05			
Nitrate (as N)	mg/L	13000	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-			
Nitrite (as N)	mg/L	60	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-			
Ammonia (as N)	mg/L	19	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05		
Total Kjeldahl Nitrogen	mg/L	-	-	0.4	0.4	0.3	0.3	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-			
Total Organic Carbon	mg/L	1.9	3.6	3	4.4	<0.5	3.7	4.6	3.6	2.7	4	2.4	4.3	2	2.2	2.4	4.3	3.3	3.1	5	4.5	2.9	2.4	4.1	3.3	2.7	3.5	4.2	<5	2.7	2.9	3.1	4				
Ortho Phosphate (as P)	mg/L	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01			
Total Phosphorus	ug/L	2.5	7	12	25	15	8	20	11	12	7	14	14	9	12	7		13	12	20	3	5	7	8	2	31	21	19	17	20	10	14	4				
Dissolved Phosphorus	mg/L	<0.005	-	<0.1	-	0.006	-	-	<0.1	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-				
pH	Units	6.5-9	7.0	7.0	7.1	7.3	7.0	6.8	7.4	7.5	7.4	7.5	7.6	7.5	7.5	7.6	7.4	7.4	7.1	7.4	7.5	7.5	7.3	7.5	7.5	7.5	7.4	7.5	6.8	7.6	7.5	7.70	7.80	7.73			
Reactive Silica (as SiO <sub>2</sub> )	mg/L	1.8	0.7	2.1	3.5	2.5	1.3	2.5	<0.5	0.9	0.7	2.3	0.8	1.8	0.8	2.6	3.6	2.2	0.6	2.2	3.4	2.8	1.2	2.4	2.4	1.9	0.8	2.0	2.5	1.9	1.0	2.0	3.5				
Sulphate	mg/L	16	12	13	13	15	15	11	12	17	18	16	13	20	24	23	20	25	26	20	17	18	16	16	23	17	16	13	16	18	15	13					
Turbidity	NTU	0.7	0.6	0.8	2	1.3	1.1	1	3.6	2.9	0.9	0.8	3.2	3.1	1	1.7	0.7	1.4	0.6	1	0.8	1.1	0.5	1	1.2	2.8	1.0	2.2	1.2	1.2	1.1	0.90	1.30				
Conductivity	µS/cm	570	360	390	310	350	420	330	340	630	560	470	390	680	910	780	700	1000	960	810	570	620	720	630	580	930	640	540	390	650	700	700	480				
TDS (calculated)	mg/L	289	176	206	159	192	212	176	170	326	285	245	204	345	462	434	368	518	485	418	300	327	358	139	309	456	332	279	207	328	365	359	256				
Bicarbonate (as CaCO <sub>3</sub> )	mg/L	20	17.1	21.3	17	15	20	25	26	22	29	27	30	19	22	27	34	27	30	36	35	30	35	36	36	31	31	33	35	27	33	35	36				
Carbonate (as CaCO <sub>3</sub> )	mg/L	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1
Calcium	mg/L	16	12	14	12	13	14	13	14	20	18	17	15	18	21	22	19	24	22	23	18	18	23	22	20	24	20	18	16	19	21.4	21	17.5				
Magnesium	mg/L	1.8	1.4	1.7	1.6</td																																

TABLE 9 Surface Water Quality Data for Russell Lake, Outlet (2005-2012)

Analyte	Units	CCME FWAL	Outlet																															
			2005				2006				2007				2008				2009				2010				2011							
			Apr	Jun	Aug	Nov	Mar	Jun	Aug	Oct	Apr	Jun	Aug	Oct	Mar	Jun	Aug	Oct	Apr	Jun	Aug	Oct	Mar	Jun	Aug	Nov	Apr	Jun	Aug	Nov				
<b>General Chemistry</b>																																		
Total Alkalinity (asCaCO <sub>3</sub> )	mg/L		19	17	23	17	16	21	25	26	22	26	27	30	19	23	27	33	27	29	36	34	30	34	37	37	29	33	34	34	28	30	37	37
Chloride	mg/L		150	88	100	72	93	99	79	80	160	140	120	90	180	250	220	190	270	260	220	150	170	180	170	150	240	170	130	93	180	190	190	120
Colour	TCU		11	12	12	18	12	14	14	11	9	9	12	11	10	16	9	16	10	16	18	16	14	18	13	21	21	22	13	8.7	10	16		
Hardness (as CaCO <sub>3</sub> )	mg/L		49	6.94	42	37	-	43	41	41	59	54	50	44	53	63	59	65	68	66	51	56	73	62	58	68	55	54	48	58	61	63	55	
Nitrate + Nitrite (as N)	mg/L	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	0.14	<0.05	<0.05	0.21	<0.05	<0.05	<0.05	<0.05	0.07	<0.05	<0.05	<0.05	0.07	<0.05	<0.05	<0.05	0.092	<0.05	<0.05	<0.05	
Nitrate (as N)	mg/L	13000	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	-	-	-	-	-	-	-	-	-	<0.05	-	-	-	-	-	-	-	-	-	-	-		
Nitrite (as N)	mg/L	60	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	-	-	-	-	-	-	<0.01	-	-	<0.01	<0.01	-	-	-	-	-	-	-	-	-
Ammonia (as N)	mg/L	19	<0.05	<0.05	0.06	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05
Total Kjeldahl Nitrogen	mg/L	-	0.5	0.4	0.3	0.1	0.4	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
Total Organic Carbon	mg/L	2.1	3.6	4.3	3.6	3.3	5.4	2.9	2.9	3.7	3.2	4.3	2.3	3.3	3.2	3.8	2.7	3.1	3.7	3.5	2.8	3	3.7	3.6	2.5	3.7	4.2	<0.01	3	3.2	3.5	4.1		
Ortho Phosphate (as P)	mg/L	0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	
Total Phosphorus	ug/L	7	10	16	27	10	4	<20	10	14	73	14	12	9	15	8	17	11	17	6	6	14	7	7	23	17	16	13	21	5	11	8		
Dissolved Phosphorus	mg/L	0.005	-	<0.1	-	0.007	-	-	<0.1	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
pH	Units	6.5-9.0	6.7	6.9	7.1	7.3	6.9	6.7	7.4	7.5	7.2	7.4	7.6	7.5	7.5	7.4	7.3	7.3	7.1	7.5	7.3	7.3	7.7	7.5	7.5	7.7	6.6	7.6	7.5	7.7	7.6	7.7		
Reactive Silica (as SiO <sub>2</sub> )	mg/L	1.4	1	2.5	3.5	2	1.3	2.4	1.4	0.9	4.4	2.3	0.8	1.8	1.6	2.6	3.4	2.1	0.5	2.4	3.3	2.3	0.9	2.4	2.4	1.7	0.8	2.1	2.4	1.3	2.4	3.4		
Sulphate	mg/L	16	12	12	16	14	14	11	12	17	18	16	13	19	24	23	20	17	17	17	18	17	16	23	17	15.0	13	16	18	15	12			
Turbidity	NTU	1.4	2.4	0.2	1.8	1.2	1	0.3	2.8	2.9	0.7	0.6	2.5	2.8	0.8	1.2	0.3	1.4	0.6	0.5	0.7	1.2	1	0.6	0.8	2.4	0.6	1.3	1.1	1.8	0.88	0.34	1.3	
Conductivity	µS/cm	580	360	400	300	360	410	320	330	620	550	470	390	690	910	780	700	1000	970	810	570	630	750	630	580	930	640	540	390	660	700	710	480	
TDS (calculated)	mg/L	292	<1	214	162	197	209	178	171	323	286	243	201	342	469	435	372	506	499	417	296	328	365	319	308	450	324	279	207	339	360	365	260	
Bicarbonate (as CaCO <sub>3</sub> )	mg/L	19.5	36	23	17	16	21	25	26	22	26	27	30	19	23	27	33	27	29	36	34	30	34	37	36	29	33	33	34	28	30	37	36	
Carbonate (as CaCO <sub>3</sub> )	mg/L	<1	16.8	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1
Calcium	mg/L	17	12	14	12	14	14	14	14	20	18	17	15	18	22	22	19	22	23	23	17	19	25	21	20	23	19	18	16	20.0	20.7	21.3	148.3	
Magnesium	mg/L	1.8	1.5	1.7	1.6	1.7	1.7	1.8	2.3	2.2	2.1	1.8	1.8	2.2	2.2	2.1	2.2	2.3	2.3	1.9	2.0	2.7	2.3	2.1	2.3	2.0	2	1.9	2.0	2.24	2.39	2.13		
Potassium																																		

**TABLE 10 Surface Water Quality Data for Russell Lake, South Inlet (2005-2012)**

---

FWAL - Freshwater Aquatic Life

**\*Results below detection limits (DL) shown as 1/2 DL**

**\*\*Results below detection limits shown as 1**

Cells left intentionally blank for graphs to indicate n

Cells with dash indicate no sample collected

TABLE 11 Surface Water Quality Data for Russell Lake, North Inlet (2005-2012)

Analyte	Units	CCME FWAL	North Inlet																																	
			2005				2006				2007				2008				2009				2010				2011									
			Apr	Jun	Aug	Nov	Mar	Jun	Aug	Oct	Apr	Jun	Aug	Oct	Mar	Jun	Aug	Oct	Apr	Jun	Aug	Oct	Mar	Jun	Aug	Nov	Apr	Jun	Aug	Nov						
<b>General Chemistry</b>																																				
Total Alkalinity (as CaCO <sub>3</sub> )	mg/L		28	-	56	-	71	63	70	41	56	81	88	83	40	<1	87	71	47	80	92	72	70	110	99	77	83	92	77	98	84	68	76	93		
Chloride	mg/L		110	-	110	-	590	130	140	87	390	200	150	150	1500	350	230	91	270	310	260	190	640	330	270	190	560	290	130	250	560	190	200	210		
Colour	TCU		28	-	17	-	7	21	13	20	12	13	15	12	7	16	49	13	30	24	25	27	11	22	38	33	13	36	63	18	9.3	41	21	20		
Hardness (as CaCO <sub>3</sub> )	mg/L		62	-	93	-	-	86	110	81	160	150	120	120	230	150	130	120	100	140	140	110	200	170	160	120	220	150	90	170	210	99	110	130		
Nitrate + Nitrite (as N)	mg/L	13000	0.65	-	0.22	-	0.4	<0.05	0.07	1.3	0.32	0.24	0.25	0.23	0.62	0.2	0.26	0.15	0.56	0.27	0.21	0.23	0.33	0.26	0.24	0.36	0.47	0.27	0.38	0.35	0.27	0.74	0.27	0.32		
Nitrate (as N)	mg/L	60	<0.01	-	-	-	<0.01	<0.01	<0.01	0.03	<0.01	-	-	-	-	-	-	-	<0.01	-	-	0.01	<0.01	-	-	-	-	-	-	-	-	-	-			
Nitrite (as N)	mg/L	19	<0.05	-	<0.05	-	0.14	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	0.12	<0.05	<0.05	<0.05	0.06	<0.05	0.06	<0.05	<0.05	0.15	0.07	<0.05	0.06	0.1	0.12	0.15	0.082	<0.05	0.091
Total Kjeldahl Nitrogen	mg/L		-	-	0.3	-	0.6	0.4	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-			
Total Organic Carbon	mg/L	6.5	-	3.3	-	2	4.6	5.8	4.9	4.2	3.9	3.7	4.3	5	3.8	6.2	3.3	5	4.7	4.8	5	3.4	5	7	4.5	3.1	5.1	6.6	6	3.3	6.2	4.4	4.5			
Ortho Phosphate (as P)	mg/L	0.02	-	<0.01	-	<0.01	<0.01	0.11	<0.01	5.2	0.03	0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	0.010	0.010	<0.01			
*Total Phosphorus	ug/L	29	-	22	-	8	9	180	10	2.5	56	23	12	180	15	6		73	11	10	7	7	8	14	6	15	27	50	27	7	15	2	9			
Dissolved Phosphorus	mg/L	-	-	-	-	0.007	-	-	-	5.1	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-				
pH	Units	6.5-9.0	7.3	-	7.6	-	7.7	7.3	7.8	7.6	7.3	7.9	7.9	7.6	8.0	7.8	7.7	7.4	7.7	7.9	7.9	7.8	7.9	7.9	7.9	7.9	8.0	8.0	7.98	8.1	8.04					
Reactive Silica (as SiO <sub>2</sub> )	mg/L	3.3	-	4	-	3.4	2.2	3.3	5.5	3.3	3.3	4.9	4.2	2.6	2.2	5.7	4.5	3.2	2.9	4.9	5.4	4.5	5.5	5.9	6.1	5.3	4.1	5.8	3.8	5.2	5	6.4				
Sulphate	mg/L	26	-	21	-	40	17	12	26	46	50	30	23	97	22	32	20	38	28	14	23	<0.01	17	19	29	46	23	22.0	22	36	28	15	19			
Turbidity	NTU	25	-	3.3	-	14	2.1	0.7	10	4.2	1.5	1.2	1.5	1.2	1.5	1.70	0.8	1.5	0.6	8.5	1	0.9	1.3	1.8	2.7	1.3	1.3	1.8	4.3	1.9	2	5.1	1.7	1.3		
Conductivity	μS/cm	490	-	520	-	2000	580	620	430	1500	900	720	670	5100	1400	890	780	1100	1300	1100	760	2200	1400	1100	790	2100	1100	610	1000	2000	800	830	890			
TDS (calculated)	mg/L	262	-	274	-	1110	310	339	234	804	488	392	364	2740	693	521	413	563	652	560	407	1170	-	574	428	1080	593	335	544	1030	427	439	468			
Bicarbonate (as CaCO <sub>3</sub> )	mg/L	28	-	56	-	71	63	70	41	56	80	88	83	40	83	87	70	47	80	91	71	70	104	98	76	82	92	76	97	84	67	75	92			
Carbonate (as CaCO <sub>3</sub> )	mg/L	<1	-	<1	-	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1			
Calcium	mg/L	21	-	31	-	62	29	39	26	55	50	42	40	78	52	43	40	35	47	48	37	69	57	54	40	75	51	31	57	71	34	39	44			
Magnesium	mg/L	2.3	-	3.7	-	6.3	3.2	4	3.7	5	5.2	4.4	4.3	8.4	5.8	4.4	4.8	3.8	5.1	5.6	4.3	7.7	7	5.8	4.2	8.6	6.1	3.3	7.3	8.3	3.7	4.4	5.4			
Potassium	mg/L	2.3	-	2.4	-	3.4	2.1	2.5	5.2	5.9	4.4	4.1	4.8	6.8	2.6	3.4	3.3	3.4	2.9	3.0	2.6	4.0	3.4	3.7	3.1	3.8	2.8	2.9	3.1	3.5	3.1	2.3	2.6			
Sodium	mg/L	75	-	68	-	360	83	91	49	250	120	98	90	210	150	100	180	200</																		