

Stantec Consulting Ltd. 102 - 40 Highfield Park Drive Dartmouth NS B3A 0A3 Tel: (902) 468-7777

Fax: (902) 468-9009

December 9, 2010 File: SD19184/121510291

Clayton Developments Limited 255 Lacewood Drive, Suite 100C Halifax, NS B3M 4G2

Attention: Mike Hanusiak, Sr. Vice President, General Manager

Dear Mr. Hanusiak:

Reference: Water Quality Monitoring Results for Russell Lake – November 2011 Sampling Event

To monitor the effects of development, water quality monitoring has been undertaken on a seasonal basis since April of 2005 at four stations in Russell Lake (Figure A1 in Attachment). Monitoring events include one spring, two summers and one fall sampling event each year. Water samples are not collected in winter. Sampling results have been provided in previous letter reports to you, with a summary of the sampling up to August 10, 2011 presented in the report dated August 30, 2011. The present report builds on the results to date while focusing on the monitoring results for the November 9, 2011 fall sampling event.

The weather was calm and cloudy with an air temperature of 4°C at the time of sampling, and cloudy for four days prior to the sampling event on November 9, 2011. Winds blew generally from the west and were below 30 km/hr for the four days leading up to the day of sample collection, and less than 10 km/hr the day of sample collection.

Sampling was conducted between 11:15 a.m. and 12:35 p.m. on November 9, 2010. At the time of sample collection, the water at the North Inlet station was observed to be opaque and brown. The grating within the culvert at this station was obstructed by garbage but no smell was noted. Water flowing from the South Inlet was clear and the flow was moderate. No algae were observed and the aquatic vegetation was limited to attached macrophytes. No sedimentation was observed in the pools. No birds were observed in the wetlands near the South Inlet. At the In-Lake station, approximately ten gulls were observed about 30 m south of the sampling location and ducks were observed along the northern shore. Algal growth was present on the dock. The water at the In-Lake station was brown and turbid. At the Outlet station of Russell Lake the water was running clear and no odour was detected. Filamentous algae covered the substrate at this station. The aquatic vegetation had died off for the year by the date of sampling.

Tables A1 - A4 (in Attachment) present the water quality data for the period from April 2005 to November 2011 at the four Russell Lake monitoring stations: In-Lake, Outlet, South Inlet and North Inlet sampling stations. The trends for some of the key parameters (total suspended solids (TSS), pH, conductivity, sodium and chloride, total phosphorus (TP), chlorophyll *a* and faecal coliforms) are presented graphically in Figures A2 to A8 (in Attachment).

December 9, 2011 Mike Hanusiak, Sr. Vice President, General Manager Page 2 of 7

Reference: Water Quality Monitoring Results for Russell Lake – November 2011 Sampling Event

Total Suspended Solids

TSS concentrations (Figure A2 in Attachment) were low with a concentration of 3 mg/L at the In-Lake station, 2 mg/L at the North Inlet and Outlet stations, and 1 mg/L at the South Inlet. These results are consistent with previous fall results. The TSS measured in the duplicate field sample for the North Inlet station was 3 mg/L, which is similar to the result for the field sample (2 mg/L).

pН

Measurements of pH collected in the field since April 2005 (Figure A3 in Attachment) indicate that the waters of Russell Lake are generally neutral. The pH readings in early November 2011 were slightly acidic at pH 5.7 for both the In-Lake and South inlet stations and which were below the CCME freshwater aquatic guideline of pH 6.5. The historic range for these two stations in the fall is slightly higher with a pH from 6.8 to 7.7 for the In-Lake station and a pH from 6.7 to 7.7 for the South Inlet station. The North Inlet station had a higher pH of 8.1 than what would normally be encountered in the fall and which would range from pH 7.5 to 7.9. The Outlet station measured pH 6.8 and within the historic range of pH 6.8 to 7.8.

Salt Concentrations

Russell Lake has higher levels of salt concentrations due to the level of development within the watershed. This is typical of any urban lake when compared to a similar lake in a less developed watershed. Conductivity increases with salt concentrations and elevated levels are typically exhibited during spring melt conditions. Concentrations tend to be higher at the North Inlet where the road network and associated drainage is concentrated and lower at the South Inlet which is less developed.

The November 2011 conductivity, chloride (CI) ion, and sodium (Na) concentrations were generally within the range of past fall sampling results. As is typical, the concentrations at the North Inlet were higher when compared to the other three stations (Tables A1 to A4; Figures A4 and A5 in Attachment).

The fall 2005 to 2010 values for the North Inlet station ranged from 430 to 790 µS/cm for conductivity, from 49 to 110 mg/L for sodium, and from 87 to 190 mg/L for chloride. In early November 2011, the concentrations measured at the North Inlet station for conductivity, sodium and chloride were 1000 µS/cm, 132 mg/L and 250 mg/L, respectively. The 2011 results at this station were all slightly higher than the range of previous results.

The values measured in the fall of 2005 to 2010 at the South Inlet station ranged from 120 to 210 μ S/cm for conductivity, with results for sodium ranging from 13 to 23 mg/L and chloride ranging from 20 to 37 mg/L. The early November 2011 values at this station were within that range with conductivity measuring at 150 μ S/cm, sodium at 16 mg/L and chloride at 27 mg/L.

The fall values for the In-Lake sampling station for 2005 to 2010 ranged from 310 to 700 μ S/cm for conductivity, from 45 mg/L to 110 mg/L for sodium and from 72 to 190 mg/L for chloride. In early November 2011 the values at this station were within that range with conductivity measuring at 390 μ S/cm, sodium at 57 mg/L and chloride at 94 mg/L.

Similar values are typically observed between the Outlet station and the In-Lake station, as supported by the range of fall values at the Outlet station for conductivity (300 to 700 μ S/cm), sodium (44 to 110 mg/L) and chloride (72 to 190 mg/L) for the fall 2005 to 2010sampling events. In early November 2011 at the Outlet

December 9, 2011 Mike Hanusiak, Sr. Vice President, General Manager Page 3 of 7

Reference: Water Quality Monitoring Results for Russell Lake – November 2011 Sampling Event

station, the results were within this range with conductivity measured at 390 μ S/cm, sodium at 58 mg/L and chloride at 93 mg/L.

Nutrient Enrichment

The chlorophyll a (Chl a) concentration (based on the acidification technique) at the In-Lake station for November 2011 was 18.67 µg/L. This concentration is higher than results for the previous fall events. The fall results for the previous sampling years ranged from 1.48 µg/L in 2008 to 16.2 µg/L in 2005 (Figure A7 in Attachment). Chl a concentrations were measured at 6.58 µg/L, 7.11 µg/L, 1.48 µg/L, 4.78 µg/L, and 3.96 µg/L in 2006, 2007, 2008, 2009, and 2010 respectively. Total phosphorus (TP) concentrations at the In-Lake site have fluctuated over the fall sampling periods, ranging from a low of 2 µg/L in 2010 to a high of 25 µg/L in November of 2005 (Figure A6 in Attachment). The November 2011 result was 17 µg/L which is within the range of previous values but above the HRM threshold value of 15 µg/L for Russell Lake. It is noted that the threshold value is intended for comparison to sample results at the In-Lake station only.

The ChI a concentration at the Outlet station measured 16.61 ug/L in November 2011. The 2011 result is within the range of the 2005 to 2010 values which ranged from 1.04 μ g/L in October 2008 to 17.7 μ g/L in November 2005 but higher than the trend since 2006 which has been less than 5.46 μ g/L for the fall results. The November 2011 TP concentration at the Outlet station measured 13 μ g/L which is within the range of previous fall results (Table A2 in Attachment) and ranged from a low of 6 μ g/L in 2009 to a high of 27 μ g/L in 2005.

The ChI a concentration for the fall sampling event at the South Inlet station was 0.13 μ g/L. This result is lower than the range observed for all previous fall values of between 0.15 μ g/L in 2007 to 1.08 μ g/L in 2006. The TP concentration at the South Inlet station in November 2011 measured 31 μ g/L and which is in the range of all past fall results that ranged from 26 μ g/L in 2010 to 88 μ g/L in 2006.

At 1.61 μ g/L, the ChI a concentration at the North Inlet station in November 2011 was within the range of previous fall results where the fall 2007, 2006, 2009, 2010 and 2008 values of 6.60 μ g/L, 1.74 μ g/L 1.44 μ g/L, 1.04 μ g/L, and 0.17 μ g/L were obtained. ChI a concentrations were not measured in the fall of 2005. The TP concentration for the North Inlet station measured 27 μ g/L which is the highest fall value obtained since 2006 (TP concentrations were not measured in the fall of 2005). The next highest TP value of 12 μ g/L in the fall was measured in 2007.

Total Phosphorus Discussion

Total phosphorus (TP) concentrations at the In-Lake site have fluctuated over the fall sampling periods, ranging from a low of 2 μ g/L in 2010 to a high of 25 μ g/L in November of 2005 (Figure A6 in Attachment). The November 2011 result was 17 μ g/L which is within the range of previous values but above the HRM threshold value of 15 μ g/L for Russell Lake. It is noted that the threshold value is intended for comparison to sample results at the In-Lake station only. TP results for the fall 2008 sampling event were invalid.

In 2011, the In-Lake TP concentrations were above threshold values for each of the four sampling events. This led to further investigations to attempt to correlate the In-Lake TP concentrations to the North or South Inlet stations. Annual mean concentrations were calculated for the In-Lake and inlet stations to aid in determining any annual trends (Table 1). The 2011 mean for the In-Lake station was the highest observed since the beginning of monitoring at 22 ug/L. The mean TP concentrations for 2011 for the North and South inlet stations were observed to be within ranges previous measured and just above the 2005 to 2011 means of 48 ug/L and 33 ug/L, respectively.

December 9, 2011 Mike Hanusiak, Sr. Vice President, General Manager Page 4 of 7

Reference: Water Quality Monitoring Results for Russell Lake – November 2011 Sampling Event

Table 1 Annual mean TP concentrations for the In-Lake and Inlet stations (2005-2011)

Year	Annual In-Lake TP Mean	Annual N-Inlet TP Mean	Annual S-Inlet TP Mean
2005	12	55	26
2006	14	77	51
2007	12	27	23
2008	9	32	67
2009	12	60	25
2010	6	41	9
2011	22	46	30
2005 - 2011	12	48	33

The Pearson correlation coefficient statistic was used on individual values and not the mean values to determine which of the inlet stations most strongly correlated to the In-Lake TP concentrations (Table 2). The Pearson correlation coefficient ranges from -1 to 1. A value of 1 implies that a linear equation describes the relationship between *Inlet* and *In-Lake concentrations* perfectly, with all data points lying on a line for which *In-Lake concentrations* increase. A value of -1 implies that all data points lie on a line for which *In-Lake concentrations* decrease as *Inlet concentrations* increase. A value of 0 implies that there is no linear correlation between the variables. This exercise in correlation was completed for the North and South Inlet stations for all years of monitoring and subsequently for their sum as total contribution of TP to Russel Lake. The correlation coefficients for comparison of Inlet stations with the In-Lake station TP concentrations were very low.

Table 2 Pearson correlation coefficients between the In-Lake and inlet stations for TP concentrations

Inlet	Pearson Correlation (2005-2011)
North Inlet TP Concentration	0.0159
South Inlet TP Concentration	0.0190
Sum of Inlet TP Concentrations	0.0215

A method of visual correlation of the Inlet stations and In-Lake station concentrations illustrate that high concentrations of TP in the Inlet stations are not strongly associated with elevated levels in the In-Lake station. Figures 1, 2 and 3 below illustrate the data with a trend line through the data illustrating the linear regression equation and the R-squared coefficient (R²). The R-squared coefficient represents how the data fits the linear regression equation. A value of 1 implies that the data completely fits the linear regression equation with a value of 0 implying that the data points do not fit the equation. The R-squared coefficients are low, varying from 0.0245 to 0.0363. Based on the results of the Pearson correlation and linear regression, the elevated TP concentrations observed for the In-Lake station cannot be correlated to the TP concentrations observed coming from the North or South Inlet stations. This suggests that there are other sources of TP within Russell Lake.

December 9, 2011 Mike Hanusiak, Sr. Vice President, General Manager Page 5 of 7

Reference: Water Quality Monitoring Results for Russell Lake – November 2011 Sampling Event




Figure 1 Correlation of North Inlet TP Concentrations to In-Lake TP Concentrations




Figure 2 Correlation of South Inlet TP Concentrations to In-Lake TP Concentrations

December 9, 2011 Mike Hanusiak, Sr. Vice President, General Manager Page 6 of 7

Reference: Water Quality Monitoring Results for Russell Lake – November 2011 Sampling Event




Figure 3 Correlation of the sum of TP Concentrations from both Inlet Stations to the In-Lake TP Concentrations

Bacterial Contamination

Bacterial contamination in Russell Lake is measured by sampling and testing for the presence of faecal coliforms in the water. In addition to faecal coliform analysis conducted by the lab, analysis of *Escherichia coli* (*E. coli*) has been added. *E coli* are a type of faecal coliform but may be considered a more specific indicator of bacteria found in the intestines of warm blooded animals and humans and a strong indication of recent sewage or animal waste contamination, whereas some faecal bacteria may originate from non-faecal sources. *E coli* counts measured 4 Colony Forming Unit (CFU)/100 mL for the In-Lake and Outlet stations, and 5 CFU/100mL for the South Inlet station (Figure A8 in Attachment). At the North Inlet station and in the duplicate field sample, *E coli* counts were 14 CFU/100 mL and 11 CFU/100 mL, respectively.

Samples analyzed for faecal coliforms at the four stations during early November 2011 measured below the detection limit (<1 Most Probable Number (MPN)/100 mL) at the In-Lake station and Outlet station, 11 MPN/100 mL at the South Inlet, 15 MPN/100 mL at the North Inlet and 25 MPN/100 mL in the duplicate sample. The geometric mean of the results for faecal coliform analysis of 4 MPN/100 mL indicates that faecal coliform levels were below the CCME Recreational Guideline (200 MPN/100 mL) for freshwater at the time of sample collection. The geometric mean for the *E. coli* results is 7 CFU/100 mL, which is also below the recreational guideline value of 200 *E. coli* per 100 mL.

Sample results at all stations for faecal coliform were within the range of previous fall results. Historical fall results at the South Inlet station range from <1 MPN/100 ML (2005 and 2006) to 280 MPN/100 mL in 2010. Results for 2007 and 2009 were 6 MPN/100 mL and 2 MPN/100 mL, respectively. At the Outlet station, previous fall results range from <1 MPN/100 mL (2005 and 2006) to 640 MPN/100 mL (2007). The In-Lake station faecal coliform value was <1 MPN/100 mL, within the range of past results from <1 MPN/100 mL (2005 and 2006) to 140 MPN/100 mL (2007). Faecal coliform results at the North Inlet station are typically higher than the other stations. The faecal coliform value for the 2011 fall sample (15 MPN/100 mL) was within the range of the fall sampling events for this station, which ranged in the past from <1 MPN/100 mL (2005) to 710 MPN/100 mL (2007).

December 9, 2011 Mike Hanusiak, Sr. Vice President, General Manager Page 7 of 7

Reference: Water Quality Monitoring Results for Russell Lake - November 2011 Sampling Event

Conclusions

The results of the November 2011 sampling event indicate that total phosphorus level at the In-Lake station was 17 μ g/L, which is higher than previous fall sampling results with the exception of 2005 when the total phosphorus result was 25 μ g/L, and is slightly above the HRM threshold value of 15 μ g/L. Based on the results of the Pearson correlation and linear regression, the elevated TP concentrations observed in the In-Lake Stations are not correlated to TP concentrations observed in the North or South Inlet stations. ChI *a* concentrations were also lower or within the range for the 2011 fall event and compared to previous years, with the exception of the In-Lake station where the concentration was slightly higher than the range at 18.67 μ g/L. It is possible that the higher TP concentration observed for the In-Lake station is attributable to the higher ChI a concentration and *in-situ* algal biomass in Russell Lake.

The November 2011 conductivity, sodium (Na) and chloride (Cl) concentrations were within the range of previous sampling events for all sites. The values at the North Inlet station have tended toward higher concentrations since 2007. Higher levels of salts would be anticipated as development progresses in and around the Lake. TSS measurements were low and pH measurements indicate that the lake water is generally neutral, but which had a basic pH of 8.1 for the North Inlet station and an acidic pH of 5.7 for both the In-Lake and South Inlet stations for the fall sampling event. The latter two stations are below the CCME freshwater aquatic pH guideline of 6.5. The geometric means of all *E. coli* and faecal coliform results were below the CCME recreational guideline of 200 MPN/100mL. The overall results for the various water quality parameters measured suggest generally consistent water quality in Russell Lake for the fall 2011 sampling event compared to previous years, except for possibly pH, total phosphorous, and ChI *a.*

Sincerely,


STANTEC CONSULTING LTD


ORIGINAL SIGNED BY


Kelley Fraser, MES, EPt Project Manager Tel: (902) 468-7777 kelley.fraser@stantec.com

cc: Cameron Deacoff, Halifax Regional Municipality


Attachments: Figures A1- A8 and Tables A1- A4







*gaps are shown where data is unavailable









Figure A2. Total Suspended Solids at four sites in Russell Lake from April 2005 to November 2011










Figure A3. pH at four sites in Russell Lake from April 2005 to November 2011







Figure A4. Conductivity at four sites in Russell Lake from April 2005 to November 2011





*gaps are shown where data are unavailable







Figure A5. Sodium and chloride concentrations at four sites in Russell Lake from April 2005 to November 2011



*gaps are shown where data are unavailable





Figure A6.Total phosphorus concentrations at four sites in Russell Lake from April 2005 to November 2011



*gaps are shown where data are unavailable





Figure A7. Chlorophyll a at four sites in Russell Lake from April 2005 to November 2011



*gaps are shown where data are unavailable




Figure A8. Faecal coliforms at four sites in Russell Lake from April 2005 to November 2011

TABLE A1 Surface Water Quality Data for Russell Lake, In-Lake (2005-2011)

		20115														Ir	n-Lake													
Analyte	Units	CCME FWAL			2005			2	006			2	007			20	08			20	09			20	10			20	011	
		IWAL	Apr	Jun	Aug	Nov	Mar	Jun	Aug	Oct	Apr	Jun	Aug	Oct	Mar	Jun	Aug	Oct	Apr	Jun	Aug	Oct	Mar	Jun	Aug	Nov	Mar	Jun	Aug	Nov
General Chemistry																													7	
Total Alkalinity (asCaCO ₃)	mg/L		20	17	21	17	15	20	25	26	22	29	27	30	19	22	28	34	27	30	36	35	30	35	36	36	32	31	33	35
Chloride	mg/L		150	87	99	72	91	100	77	79	170	140	120	89	180	250	220	190	270	250	220	150	170	180	160	150	240	170	130	94
Colour	TCU		12	12	8	18	12	14	13	11	9	9	8	12	10	11	15	9	16	9	14	20	14	8	10	15	14	19	22	21
Hardness (as CaCO ₃)	mg/L		48	34	40	37		43	39	42	59	53	52	46	52	62	64	56	70	65	66	52	54	69	65	58	69	60	53	47
Nitrate + Nitrite (as N)	mg/L		<0.05	<0.05	< 0.05	< 0.05	< 0.05	<0.05	<0.05	<0.05	< 0.05	<0.05	<0.05	<0.05	0.15	<0.05	<0.05	<0.05	0.2	< 0.05	<0.05	< 0.05	0.11	< 0.05	<0.05	< 0.05	0.08	< 0.05	<0.05	<0.05
Nitrate (as N)	mg/L	13000	<0.05	<0.05	< 0.05	< 0.05	< 0.05	<0.05	<0.05	<0.05	<0.05	-	-	=	-	=	-	-	-	-	-	<0.05	-	-	<0.05	< 0.05	-	-	-	-
Nitrite (as N)	mg/L	60	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	-	-	-	-	-	-	-	-	-	-	<0.01	1	-	<0.01	<0.01	-	-	-	-
Ammonia (as N)	mg/L	19	<0.05	<0.05	< 0.05	< 0.05	< 0.05	<0.05	<0.05	<0.05	>0.05	<0.05	<0.05	<0.05	< 0.05	<0.05	< 0.05	<0.05	< 0.05	< 0.05	<0.05	< 0.05	< 0.05	< 0.05	<0.05	<0.05	< 0.05	< 0.05	0.12	<0.05
Total Keljdahl Nitrogen	mg/L		-	-	0.4	0.4	0.3	0.3	-	-	-	-	-	=	-	=	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Total Organic Carbon	mg/L		1.9	3.6	3	4.4	<0.5	3.7	4.6	3.6	2.7	4	2.4	4.3	2	2.2	2.4	4.3	3.3	3.1	5	4.5	2.9	2.4	4.1	3.3	2.7	3.5	4.2	<5
Ortho Phosphate (as P)	mg/L		<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01
Total Phosphorus	ug/L		2.5	7	12	25	15	8	20	11	12	7	14	14	9	12	7		13	12	20	3	5	7	8	2	31	21	19	17
Dissolved Phosphorus	mg/L		<0.005	-	<0.1	=	0.006	-	-	-	<0.1	-	-	-	-	-	-	-	-	-	-	-	1	-	-	-	-	-	-	-
рН	Units	6.5-9	7.0	7.0	7.1	7.3	7.0	6.8	7.4	7.5	7.4	7.5	7.6	7.5	7.5	7.6	7.4	7.4	7.1	7.4	7.5	7.5	7.3	7.5	7.5	7.5	7.4	7.5	6.8	7.6
Reactive Silica (as SiO ₂)	mg/L		1.8	0.7	2.1	3.5	2.5	1.3	2.5	<0.5	0.9	0.7	2.3	0.8	1.8	0.8	2.6	3.6	2.2	0.6	2.2	3.4	2.8	1.2	2.4	2.4	1.9	0.8	2.0	2.5
Sulphate	mg/L		16	12	13	13	15	15	11	12	17	18	16	13	20	24	23	20	25	26	20	17	18	18	16	16	23	17	16	13
Turbidity	NTU		0.7	0.6	0.8	2	1.3	1.1	1	3.6	2.9	0.9	0.8	3.2	3.1	1	1.7	0.7	1.4	0.6	1	8.0	1.1	0.5	1	1.2	2.8	1.0	2.2	1.2
Conductivity	μS/cm		570	360	390	310	350	420	330	340	630	560	470	390	680	910	780	700	1000	960	810	570	620	720	630	580	930	640	540	390
TDS (calculated)	mg/L		289	176	206	159	192	212	176	170	326	285	245	204	345	462	434	368	518	485	418	300	327	358	139	309	456	332	279	207
Bicarbonate (as CaCO ₃)	mg/L		20	17.1	21.3	17	15	20	25	26	22	29	27	30	19	22	27	34	27	30	36	35	30	35	36	36	31	31	33	35
Carbonate (as CaCO ₃)	mg/L		<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1
Calcium	mg/L		16	12	14	12	13	14	13	14	20	18	17	15	18	21	22	19	24	22	23	18	18	23	22	20	24	20	18	16
Magnesium	mg/L		1.8	1.4	1.7	1.6	1.7	1.8	1.8	1.8	2.5	2.1	2.1	1.8	1.8	2.2	2.2	2.1	2.4	2.2	2.3	2	1.9	2.6	2.4	2.2	2.3	2.2	2.0	1.9
Potassium	mg/L		1.5	1.2	1.5	1.5	1.5	1.4	1.6	1.7	2.2	2.3	2.1	2.3	2.1	2.3	2.4	2.4	2.6	2.5	2.5	2.3	2.0	2.2	2.1	2.3	2	1.9	1.8	1.9
Sodium	mg/L		92	52	64	46	58	64	54	45	100	85	73	63	110	150	140	110	180	160	130	89	95	107	89	94	144	102	85	57
*Total Suspended Solids	mg/L		1	<u>1</u>	1	<u>1</u>	<u>1</u>	3	1	4	4	1	1	4	9	2	2	1	3	1	3	1	<u>1</u>	<u>1</u>	2	1	3	2	3	3
Chlorophyll a	μg/L			1.60	2.80	16.20		1.42	4.35	6.58	9.69	1.07	3.45	7.11	1.44	1.47	3.64	1.48	6.43	1.17	26.80	4.78	1.99	0.96	2.99	3.96	29.01	3.04	8.11	18.67
Field Measurements																													1	
Secchi disc depth	m		-	3.4	2.8	3.0	-	2.8	-	1.5	-	4.8	2.5	1.5	-	-	1.75	4	1.3	6	1.4	3.75	3.5	4	2.3	2.6	1.7	1.6	1.5	1.4
Dissolved Oxygen	mg/L		10.5	8.9	7.3	13.2	-	10.6	5.6	12.3	14.5	9.8	11.0	10.2	16.9	9.5	8.0	10.8	9.7	11.3	8.8	10.9	13.4	8.1	7.6	13.1	13.1	7.4	5.9	10.5
рН	units		7.0	7.3	7.6	7.2		6.7	6.8	7.2	6.6	7.4	8.0	6.8	6.5	7.7	6.8	7.3	7.3	8.0	7.7	7.0	7.8	6.8	6.9	7.5	7.4	7.4	8.2	5.7
Conductivity	μS/cm		602	334	188	129		393	310	330	612	527	474	278	444	1701	1514	600	1031	278	831	-	601	698	635	570	917	631	526	387
Temperature (field)	°C		10.4	19.6	23.4	7.2	-	19.8	21.1	8.4	7.8	20.4	23.3	11.1	5.6	18.3	21.9	9.5	5.3	16.9	22.2	8.3	5.2	20.6	22.6	8.7	3.4	18.3	20.7	8.0
Bacteriological																														
**E.Coli	CFU/100ml																						1	<u>1</u>	3	3	<1	3	67	4
**Faecal Coliform	MPN/100 ml		<u>1</u>	<u>1</u>	1	1		5	1300	<u>1</u>	3	120	300	140	11	8	440	2	150	<u>1</u>	3200	32	1	160	20	57	2	3	240	<u>1</u>
EMAL Freehweter Aquetio		<u> </u>																												

*Results below detection limits (DL) shown as 1

Cells left intentionally blank for graphs to indicate no sample collected

Cells with dash indicate no sample collected

April 2005 result for TP is italized because result was below detection limit (DL) and shown as 2.5 (1/2 DL). The DL is now 2 ug/L.

TABLE A2 Surface Water Quality Data for Russell Lake, Outlet (2005-2011)

				,	<u>, </u>											(Outlet													
Analyte	Units	CCME FWAL			2005			2	006			2	007			20	008			20	009			2	010			21	11	
		FWAL	Apr	Jun	Aug	Nov	Mar	Jun	Aug	Oct	Apr	Jun	Aug	Oct	Mar	Jun	Aug	Oct	Apr	Jun	Aug	Oct	Mar	Jun	Aug	Nov	Mar	Jun	Aug	Nov
General Chemistry																														
Total Alkalinity (asCaCO ₃)	mg/L		19	17	23	17	16	21	25	26	22	26	27	30	19	23	27	33	27	29	36	34	30	34	37	37	29	33	34	34
Chloride	mg/L		150	88	100	72	93	99	79	80	160	140	120	90	180	250	220	190	270	260	220	150	170	180	170	150	240	170	130	93
Colour	TCU		11	12	12	18	12	14	14	11	9	9	9	12	11	10	16	9	16	10	16	18	16	14	14	18	13	21	21	22
Hardness (as CaCO ₃)	mg/L		49	6.94	42	37	-	43	41	41	59	54	50	44	53	63	65	59	65	68	66	51	56	73	62	58	68	55	54	48
Nitrate + Nitrite (as N)	mg/L		<0.05	< 0.05	<0.05	<0.05	< 0.05	<0.05	<0.05	<0.05	< 0.05	<0.05	<0.05	< 0.05	0.14	<0.05	<0.05	<0.05	0.21	<0.05	<0.05	<0.05	0.07	<0.05	<0.05	<0.05	0.07	< 0.05	<0.05	<0.05
Nitrate (as N)	mg/L	13000	< 0.05	< 0.05	<0.05	<0.05	< 0.05	<0.05	<0.05	<0.05	< 0.05	-	-	-	-	-	-	-	-	-	-	<0.05	-	-	<0.05	<0.05	-	-	-	-
Nitrite (as N)	mg/L	60	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	-	-	-	-	-	-	-	-	-	-	<0.01	-	-	<0.01	<0.01	-	-	-	-
Ammonia (as N)	mg/L	19	<0.05	<0.05	0.06	<0.05	< 0.05	<0.05	<0.05	<0.05	< 0.05	<0.05	<0.05	< 0.05	< 0.05	<0.05	<0.05	<0.05	< 0.05	<0.05	<0.05	<0.05	< 0.05	<0.05	<0.05	<0.05	< 0.05	< 0.05	0.09	<0.05
Total Keljdahl Nitrogen	mg/L		-	0.5	0.4	0.3	0.1	0.4	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Total Organic Carbon	mg/L		2.1		3.6	4.3	3.6	3.3	5.4	2.9	2.9	3.7	3.2	4.3	2.3	3.3	3.2	3.8	2.7	3.1	3.7	3.5	2.8	3	3.7	3.6	2.5	3.7	4.2	<0.01
Ortho Phosphate (as P)	mg/L		0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01
Total Phosphorus	ug/L		7	10	16	27	10	4	<20	10	14	73	14	12	9	15	8		17	11	17	6	6	14	7	7	23	17	16	13
Dissolved Phosphorus	mg/L		0.005	-	<0.1	-	0.007	-	-	-	<0.1	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
ρΗ	Units	6.5-9.0	6.7	6.9	7.1	7.3	6.9	6.7	7.4	7.5	7.2	7.4	7.6	7.5	7.5	7.4	7.3	7.3	7.1	7.5	7.3	7.8	7.3	7.3	7.7	7.5	7.5	7.7	6.6	7.6
Reactive Silica (as SiO ₂)	mg/L		1.4	1	2.5	3.5	2	1.3	2.4	1.4	0.9	4.4	2.3	0.8	1.8	1.6	2.6	3.4	2.1	0.5	2.4	3.3	2.3	0.9	2.4	2.4	1.7	0.8	2.1	2.4
Sulphate	mg/L		16	12	12	16	14	14	11	12	17	18	16	13	19	24	23	20	25	24	20	17	17	18	17	16	23	17	15.0	13
Turbidity	NTU		1.4	2.4	0.2	1.8	1.2	1	0.3	2.8	2.9	0.7	0.6	2.5	2.8	0.8	1.2	0.3	1.4	0.6	0.5	0.7	1.2	1	0.6	0.8	2.4	0.6	1.3	1.1
Conductivity	μS/cm		580	360	400	300	360	410	320	330	620	550	470	390	690	910	780	700	1000	970	810	570	630	750	630	580	930	640	540	390
TDS (calculated)	mg/L		292	<1	214	162	197	209	178	171	323	286	243	201	342	469	435	372	506	499	417	296	328	365	319	308	450	324	279	207
Bicarbonate (as CaCO ₃)	mg/L		19.5	36	23	17	16	21	25	26	22	26	27	30	19	23	27	33	27	29	36	34	30	34	37	36	29	33	33	34
Carbonate (as CaCO ₃)	mg/L		<1	16.8	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1
Calcium	mg/L		17	12	14	12	14	14	14	14	20	18	17	15	18	22	22	19	22	23	23	17	19	25	21	20	23	19	18	16
Magnesium	mg/L		1.8	1.5	1.7	1.6	1.7	1.7	1.7	1.8	2.3	2.2	2.1	1.8	1.8	2.2	2.2	2.1	2.2	2.3	2.3	1.9	2.0	2.7	2.3	2.1	2.3	2.0	2	1.9
Potassium	mg/L		1.6	1.3	1.2	1.5	1.5	1.4	1.5	1.7	2.4	2.1	2	2.3	2.1	2.3	2.4	2.4	2.4	2.6	2.5	2.1	2.0	2.2	2.0	2.2	2.0	1.8	1.8	1.9
Sodium	mg/L		92	57	65	46	61	64	52	44	110	87	71	61	110	150	140	110	160	170	130	88	98	112	85	93	143	95.1	87	58
**Total Suspended Solids	mg/L		3	2.4	3.6	<u>1</u>	5	2	<u>1</u>	3	3	1	<u>1</u>	3	1	1	<u>1</u>	2	3	1	1	1	1	<u>1</u>	<u>1</u>	1	2	2	3	2
Chlorophyll a	μg/L					17.70		2.19	2.46	5.46	7.85	0.95	1.08	5.16	1.54	1.20	2.16	1.04	6.35	1.07	11.80	4.72	2.92	1.00	1.35	3.18	19.56	1.49	7.22	16.61
Field Measurements																														
Secchi disc depth	m			-	ı		-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Dissolved Oxygen	mg/L		10.2	7.4	5.5	10.8	-	11.3	8.9	11.9	14.8	8.4	11.3	10.1	19.3	8.4	7.4	10.9	9.6	6.9	8.1	10.6	11.1	5.0	5.6	13.4	12.6	7.3	5.5	11.2
oH	units		6.9	6.8	6.8	7.2		7.2	7.1	7.4	7.0	7.0	7.8	7.4	6.5	7.2	7.6	7.8	7.4	7.2	7.6	7.8	7.5	7.7	7.4	7.2	7.7	7.6	7.6	6.8
Conductivity	μS/cm		635	345	201	136		389	399	344	613	532	473	272	448	1703	1505	602	1023	278	834	-	592	702	633	571	914	634	529	389
Temperature (field)	°C		12.2	22.5	25.5	7.6	-	21.3	23.1	9.1	9.6	22.9	24.5	9.9	5.5	18.6	23.2	7.8	6.2	19.6	23.5	8.8	5.1	20.5	22.8	7.9	5.0	21.0	21	9.3
Bacteriological																														
**E.Coli	CFU/100ml						1																3	1	29	10	1	28	54	4
*Faecal Coliform	MPN/100 ml		1	8	10	1		170	2700	1	1	700	3000	640	9	160	930	4	140	24	8100	160	1	35	600	200	4	23	490	1

*Results below detection limits (DL) shown as 2.5 (1/2 DL)

*Results below detection limits shown as 1

Cells left intentionally blank for graphs to indicate no sample collected

Cells with dash indicate no sample collected

TABLE A3 Surface Water Quality Data for Russell Lake, South Inlet (2005-2011)

					<u> </u>												South Inlet														
Analyte	Units	CCME FWAL			2005				2006				200)7			20	08			20	09			20	010			20)11	
		FWAL	Apr	Jun	Aug	Nov	Mar	Jun	Aug	Sep	Oct	Apr	Jun	Aug	Oct	Mar	Jun	Aug	Oct	Apr	Jun	Aug	Oct	Mar	Jun	Aug	Nov	Mar	Jun	Aug	Nov
General Chemistry										additional																					
Total Alkalinity (asCaCO ₃)	mg/L		11	31	33	7	21	13	<5	35	13	11	25	24	25	18	23	37	32	16	34	54	24	25	41	56	27	25	35	24	28
Chloride	mg/L		25	17	21	21	23	13	9	36	26	35	27	18	20	76	24	29	37	38	49	39	23	62	31	49	37	71	43	28	27
Colour	TCU		24	27	21	45	14	68	380	44	38	23	47	52	41	17	52	150	51	51	50	150	57	26	47	87	59	22	83	200	42
Hardness (as CaCO ₃)	mg/L		24	40	45	23	-	21	46	48	29	30	33	33	28	43	29	41	37	27	40	53	32	42	47	54	43	42	40	31	38
Nitrate + Nitrite (as N)	mg/L		<0.05	<0.05	0.09	0.08	0.13	<0.05	<0.05	0.12	0.08	0.09	<0.05	0.07	0.05	0.1	<0.05	0.1	<0.05	0.15	<0.05	0.14	<0.05	0.11	0.06	0.1	<0.05	<0.05	<0.05	<0.05	42
Nitrate (as N)	mg/L	13000	<0.05	<0.05	0.09	0.08	0.13	<0.05	<0.05	-	0.07	0.09	-	-	-	-	-	-	-	-	-	-	<0.05	-	-	0.1	<0.05	-	-	-	-
Nitrite (as N)	mg/L	60	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	-	<0.01	<0.01	-	-	-	-	-	-	-	-	-	-	<0.01	-	-	<0.01	0.01	-	-	-	-
Ammonia (as N)	mg/L	19	<0.05	<0.05	<0.05	<0.05	0.05	<0.05	0.07	< 0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	0.06	<0.05	<0.05	<0.05	0.08	<0.05	< 0.05	<0.05	0.06	<0.05	<0.05	< 0.05	0.06	<0.05
Total Keljdahl Nitrogen	mg/L		-	0.3	0.4	0.3	0.3	0.5	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Total Organic Carbon	mg/L		6.6	7.1	6.1	9	4.6	<0.5	45	11	7.2	5.6	11	11	9.2	3.3	8.5	17	11	6.3	8.9	13	7.8	5.5	8.1	12	8.4	4.2	9.1	17	5
Ortho Phosphate (as P)	mg/L		0.04	0.03	0.04	0.08	0.02	0.08	0.01	0.11	0.05	<0.01	0.05	0.07	0.03	<0.01	0.03	0.03	0.02	0.01	0.01	0.03	<0.01	0.01	0.03	0.02	0.03	0.01	0.02	0.03	0.01
Total Phosphorus	ug/L		42	46	43	88	31	86	80	110	53	34	10	14	44	18	36	28		30	110	71	27	45	47	46	26	30	52	70	31
Dissolved Phosphorus	mg/L		0.037	-	<0.1	-	0.021	-	-	-	-	<0.1	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
рН	Units	6.5-9.0	6.8	7.0	7.2	6.6	7.0	6.6	4.7	7.5	6.9	7.0	7.3	7.2	7.2	7.4	7.3	7.2	7.4	6.9	7.4	7.4	7.4	7.3	7.6	7.6	7.4	7.4	7.5	7.0	7.4
Reactive Silica (as SiO ₂)	mg/L		4.1	7.2	9.9	5.7	7.4	3.8	5.9	10	5.8	3.5	0.7	6.1	6.5	3.8	4.8	6.5	8.8	3.2	3.5	7.1	5.5	5.6	6.8	7.2	6.4	4.8	4.1	4.8	6.8
Sulphate	mg/L		12	5.1	7.9	16	11	<2	<10	<2	13	12	<2	<2	<2	15	2	<2	<2	15	3	<2	12	12	3	<2	18	14	<2	<2	11
Turbidity	NTU		0.3	0.2	<0.1	0.7	1.2	0.4	3.2	1.3	2.8	0.8	0.5	1.6	1.2	0.5	0.5	2.4	1.6	2.4	1.1	6.4	1.5	1.1	1.7	4.1	1.5	0.9	1.0	2.5	4.5
Conductivity	μS/cm		130	130	150	120	140	89	45	190	140	180	160	120	120	330	140	170	190	190	240	190	140	270	190	260	210	320	220	150	150
TDS (calculated)	mg/L		73	74.6	89.3	70	83	43	60	113	78	91	78	66	67	170	75	98	102	104	125	128	84	147	101	139	119	165	112	83	93
Bicarbonate (as CaCO ₃)	mg/L		10.7	31.3	32.8	7	21	13	<1	35	13	11	25	24	25	18	23	37	32	16	34	54	24	25	40	56	27	25	35	24	28
Carbonate (as CaCO ₃)	mg/L		<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1
Calcium	mg/L		6.6	10	11	6.7	9.1	5.8	14	14	8.4	8.3	9.4	9.3	7.9	13	8.1	13	10	8.3	12	16	9.4	12	13	16.1	13	13	12	9.7	11
Magnesium	mg/L		1.9	3.4	4.1	1.6	3.2	1.5	3	3.4	2	2.1	0.01	2.3	2	2.3	2.2	2.4	2.7	1.5	2.3	3.3	2	2.8	3.6	3.3	2.6	2.7	2.5	1.75	2.5
Potassium	mg/L		1	0.9	0.9	1	1	0.5	2	2	2.1	1.4	0.9	1	1.5	1.6	1.1	1.9	2.4	1.4	1.3	1.4	1.1	1.3	0.9	1.4	1.6	1.1	0.8	1.1	1.4
Sodium	mg/L		16	12	14	13	15	10	21	25	13	21	18	14	13	47	18	21	21	26	34	24	16	35	19	26	24	44	28	21	16
**Total Suspended Solids	mg/L		1	<u>1</u>	<u>1</u>	2	<u>1</u>	4	91	2	<u>1</u>	<u>1</u>	1	3	1	<u>1</u>	2	1	2	<u>1</u>	<u>1</u>	7	<u>1</u>	2	4	1	<u>1</u>	<u>1</u>	1	<2	5
Chlorophyll a	μg/L				0.10	0.20		1.23	2.07		1.08	2.22	0.22	0.16	0.15	1.54	0.56	0.70	0.48	0.69	1.67	2.32	0.29	1.37	5.05	0.64	0.13	1.80	0.42	1.18	0.36
Field Measurements																															
Secchi disc depth	m		-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Dissolved Oxygen	mg/L		10.4	9.3	7.4	13.0	-	10.3	8.8	-	12.3	14.8	9.9	11.7	12.5	21.4	10.5	6.9	11.4	10.0	10.9	9.1	10.7	15.6	9.3	9.5	15.1	13.7	7.9	6.3	11.0
рН	units		6.5	6.9	7.6	7.3		6.1	6.7	7.2	7.2	6.4	7.0	7.7	6.7	6.5	7.1	7.4	7.4	7.4	7.2	7.1	7.3	7.8	7.4	7.3	7.6	7.8	7.1	8.0	5.7
Conductivity	μS/cm		124	105	185	78		94	161	-	142	174	145	121	74	187	260	331	164	191	69	232	-	268	187	270	205	316	214	145.0	157.0
Temperature (field)	°C		9.0	15.3	23.5	5.7	-	14.7	16.9	-	6.6	4.8	14.7	16.3	5.1	0.5	12.0	17.2	5.6	4.3	13.3	16.0	5.7	3.7	13.0	16.6	5.4	2.1	13.3	17	7.0
Bacteriological																															
l 	CFU/100ml																							2	1 1	500	7	1	4	>250	5
**Faecal Coliform	MPN/100 ml		<u>1</u>	14	66	<u>1</u>		1	100		<u>1</u>	<u>1</u>	2	200	6	2	2	150	3	3	1	2200	2	1	1	1100	280	5	12	200	11

*Results below detection limits (DL) shown as 2.5 (1/2 DL)

*Results below detection limits shown as 1

Cells left intentionally blank for graphs to indicate no sample collected

Cells with dash indicate no sample collected

TABLE A4 Surface Water Quality Data for Russell Lake, North Inlet (2005-2011)

																Nor	th Inlet													
Analyte	Units	CCME FWAL		2	005			20	006			20	07			20	008			20	09			20)10			20	11	
		FVVAL	Apr	Jun	Aug	Nov	Mar	Jun	Aug	Oct	Apr	Jun	Aug	Oct	Mar	Jun	Aug	Oct	Apr	Jun	Aug	Oct	Mar	Jun	Aug	Nov	Mar	Jun	Aug	Nov
General Chemistry																														
Total Alkalinity (asCaCO ₃)	mg/L		28	-	56	-	71	63	70	41	56	81	88	83	40	<1	87	71	47	80	92	72	70	110	99	77	83	92	77	98
Chloride	mg/L		110		110		590	130	140	87	390	200	150	150	1500	350	230	91	270	310	260	190	640	330	270	190	560	290	130	250
Colour	TCU		28	-	17	-	7	21	13	20	12	13	15	12	7	16	49	13	30	24	25	27	11	22	38	33	13	36	63	18
Hardness (as CaCO ₃)	mg/L		62	-	93	-	-	86	110	81	160	150	120	120	230	150	130	120	100	140	140	110	200	170	160	120	220	150	90	170
Nitrate + Nitrite (as N)	mg/L		0.65	-	0.22	-	0.4	< 0.05	0.07	1.3	0.32	0.24	0.25	0.23	0.62	0.2	0.26	0.15	0.56	0.27	0.21	0.23	0.33	0.26	0.24	0.36	0.47	0.27	0.38	0.35
Nitrate (as N)	mg/L	13000		-		-	0.4	<0.05	0.07	1.27	0.32	-	=	-	=	-	-	-	-	-	-	0.23	-	-	0.23	0.36	-	-	<u> </u>	-
Nitrite (as N)	mg/L	60	<0.01	-		-	<0.01	<0.01	<0.01	0.03	<0.01	-	-	-	-	-	-	-	-	-	-	<0.01	-	-	0.01	<0.01	-	=	<u> </u>	-
Ammonia (as N)	mg/L	19	<0.05	-	<0.05	-	0.14	<0.05	<0.05	<0.05	< 0.05	< 0.05	<0.05	<0.05	0.12	<0.05	< 0.05	<0.05	0.06	< 0.05	0.06	<0.05	< 0.05	<0.05	0.15	0.07	< 0.05	0.06	0.1	0.12
Total Keljdahl Nitrogen	mg/L			-	0.3	-	0.6	0.4	•	-	=	-	-	-	=	-	-	-	-	-	-	i	=	-	-	-	-	=		=
Total Organic Carbon	mg/L		6.5	-	3.3	-	2	4.6	5.8	4.9	4.2	3.9	3.7	4.3	5	3.8	6.2	3.3	5	4.7	4.8	5	34	5	7	4.5	3.1	5.1	6.6	6
Ortho Phosphate (as P)	mg/L		0.02	-	<0.01	-	<0.01	<0.01	0.11	<0.01	5.2	0.03	0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	0.02	<0.01
*Total Phosphorus	ug/L		29		22		8	9	180	10	<u>2.5</u>	56	23	12	180	15	6		73	11	10	7	7	8	14	6	15	27	50	27
Dissolved Phosphorus	mg/L		-	-	-	-	0.007	-	•	-	5.1	-	-	-	-	-	-	-	-	-	-	ī	-	-	-	-	-	-	-]	-
рН	Units	6.5-9.0	7.3		7.6		7.7	7.3	7.8	7.6	7.3	7.9	7.9	7.9	7.6	8.0	7.8	7.7	7.4	7.7	7.9	7.9	7.8	7.9	8.0	7.8	7.9	7.9	7.9	8.0
Reactive Silica (as SiO ₂)	mg/L		3.3	-	4	-	3.4	2.2	3.3	5.5	3.3	3.3	4.9	4.2	2.6	2.2	5.7	4.5	3.2	2.9	4.9	5.4	4.5	5.5	5.9	6.1	5.3	4.1	5.8	5.8
Sulphate	mg/L		26	-	21	-	40	17	12	26	46	50	30	23	97	22	32	20	38	28	14	23	<0.01	17	19	29	46	23	22.0	22
Turbidity	NTU		25	-	3.3	-	14	2.1	0.7	10	4.2	1.5	1.2	1.5	170	0.8	1.5	0.6	8.5	1	0.9	1.3	1.8	0.9	2.7	1.3	1.3	1.8	4.3	1.9
Conductivity	μS/cm		490		520		2000	580	620	430	1500	900	720	670	5100	1400	890	780	1100	1300	1100	760	2200	1400	1100	790	2100	1100	610	1000
TDS (calculated)	mg/L		262	-	274	-	1110	310	339	234	804	488	392	364	2740	693	521	413	563	652	560	407	1170	-	574	428	1080	593	335	544
Bicarbonate (as CaCO ₃)	mg/L		28	-	56	-	71	63	70	41	56	80	88	83	40	83	87	70	47	80	91	71	70	104	98	76	82	92	76	97
Carbonate (as CaCO ₃)	mg/L		<1	-	<1	-	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1
Calcium	mg/L		21	-	31	-	62	29	39	26	55	50	42	40	78	52	43	40	35	47	48	37	69	57	54	40	75	51	31	57
Magnesium	mg/L		2.3	-	3.7	-	6.3	3.2	4	3.7	5	5.2	4.4	4.3	8.4	5.8	4.4	4.8	3.8	5.1	5.6	4.3	7.7	7	5.8	4.2	8.6	6.1	3.3	7.3
Potassium	mg/L		2.3	-	2.4	-	3.4	2.1	2.5	5.2	5.9	4.4	4.1	4.8	6.8	2.6	3.4	3.3	3.4	2.9	3.0	2.6	4.0	3.4	3.7	3.1	3.8	2.8	2.9	3.1
Sodium	mg/L		75		68		360	83	91	49	250	120	98	90	990	210	150	100	180	200	160	110	350	174	148	109	330	162	89	132
**Total Suspended Solids	mg/L		4		3		11	5	<u>1</u>	3	3	3	1	1	330	<u>1</u>	<u>1</u>	<u>1</u>	72	<u>1</u>	2	3	1	7	<u>1</u>	1	2	1	2	2
Chlorophyll a	μg/L				2.50			1.35	0.94	1.74	1.08	0.59	0.23	6.60	0.31	1.00	0.21	0.17	1.03	0.67	0.27	1.44	0.83	0.75	0.35	1.04	0.94	0.51	0.84	1.61
Field Measurements				-		-																								
Secchi disc depth	m		-	-	-	-	-	-	•	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	=		-
Dissolved Oxygen	mg/L		-	-	12.0	-	-	11.6	9.8	12.1	13.9	10.0	11.3	13.9	20.3	10.3	9.4	11.6	10.3	9.5	9.6	11.0	12.4	8.8	7.9	14.2	13.9	5.4	7.0	11.0
рН	units				7.5			7.5	7.7	7.3	7.0	7.7	7.9	7.0	7.0	8.0	7.9	7.6	8.1	7.9	7.9	7.8	8.1	7.5	7.3	7.3	7.6	7.6	7.2	8.1
Conductivity	μS/cm				170			542	754	437	1523	908	728	444	3036	2540	1750	681	1075	366	1069	-	2093	1293	1119	780	2142	1142	598	999
Temperature (field)	°C		-	_	7.2	-	-	18.9	21.8	8.2	8.9	19.0	18.7	9.0	1.9	14.4	18.3	8.7	6.4	14.8	18.3	7.5	4.1	15.8	19.0	7.2	2.5	16.1	18	8.2
Bacteriological																														
E.Coli	CFU/100ml																						45	1	21	7	1	6	>250	14
**Faecal Coliform	MPN/100 ml				<u>1</u>			35	4200	1	6	34	10000	710	22	22	830	5	6400	20	1700	51	1	65	1100	180	13	150	490	15
FWAL - Freshwater Aquatic		4 1	- 1			-	-	•			_	•	•	•	•	-		•	-	•	•		-	•	•	•				

Cells with dash indicate no sample collected

^{*}Results below detection limits (DL = 5) shown as 2.5 ug/L (1/2 DL)

^{**}Results below detection limits (DL) shown as 1

Cells left intentionally blank for graphs to indicate no sample collected